

## Plan Overview

---

*A Data Management Plan created using DMP Tool*

**Title:** Molecular and functional study of the movement of urea, water and ammonia ( $\text{NH}_3/\text{NH}_4^+$ ) through the urea transporters expressed in *Lithobates catesbeianus* oocytes

**Creator:** Raif Musa-Aziz

**Affiliation:** Universidade de São Paulo ([www5.usp.br](http://www5.usp.br))

**Data Manager:** Jessica Kabutomori Rosa, Neydiana Belize de Pina-Lopes

**Funder:** São Paulo Research Foundation ([fapesp.br](http://fapesp.br))

**Funding opportunity number:** 2018/22855-1

**Template:** Digital Curation Centre

### Project abstract:

Tissues throughout the body generate ammonia as a result of protein degradation. Upon entering circulation, this ammonia travels to the liver, where it is converted into less toxic nitrogenous compounds, urea and glutamine, which can then enter the circulation and travel to the kidneys to be excreted in the urine. In the kidney, urea contributes to maintain a high osmolarity gradient in the medullary interstitium-which is important for the urinary concentrating mechanism-and glutamine is used by the proximal tubule cells to regenerate ammonia in a metabolic process that generates equimolar concentration of new  $\text{HCO}_3^-$ -which is important to maintain acid-base homeostasis. Ammonia, as  $\text{NH}_4^+$ , is then secreted into the proximal fluid, reabsorbed by the thick ascending limb of the Henle's loop, and secreted by the collecting duct (CD) through parallel  $\text{NH}_3$  and  $\text{H}^+$  transport, which combine and re-form  $\text{NH}_4^+$  that will then be excreted. Recent observations, from our laboratory and others, indicated that the basolateral and apical membranes of CD have low  $\text{NH}_3$  permeability, suggesting that membrane proteins are likely involved in  $\text{NH}_3$  (dissolved gas) transport in the CD cell. Mammalian urea transporters (UTs) belong to the SLC14 family of solute carriers and are responsible for the facilitated diffusion of urea across plasma membrane. There are two UT genes: SLC14A1, which encodes UT-B, and SLC14A2, which generates splice variants UT-A1, A2 and A3. UT-B is highly expressed in the red blood cell membrane, and is also localized in endothelial cells of the descending vasa recta and liver. UT-A1 and A3 are expressed, respectively, on the apical and basolateral membranes of the inner medullary CD. UT-A2 is expressed in the thin descending limb of the Henle's loop. These transporters recycle and thereby concentrate urea in the renal medulla to maintain hyperosmolar interstitium, which is necessary for maximal urine

concentration, and also allow the excretion of urea with a minimal volume of water. However, there is some controversy over whether or not UTs are also physiologically relevant for H<sub>2</sub>O (during dehydration) and NH<sub>3</sub> (during acidosis) transport in the kidney. Previous work has shown that mammalian UT-B, heterologously expressed in *Xenopus laevis* oocytes, not only transport urea but are also capable of transporting H<sub>2</sub>O and NH<sub>3</sub>. The crystal structure of the bovine UT-B show that UTs are homotrimeric, proteins similar to the homotetrameric structure of AQP1. Each monomer consists of two protomers (6 helices each) that fold together to form a hydrophilic urea channel. The N- and C-termini of each monomer are located in the cytosol. At the center of the three monomers is a hydrophobic pore, blocked by lipid molecules. We hypothesize that UT-As can, in addition to urea, transport H<sub>2</sub>O and NH<sub>3</sub>, using the monomeric urea channel. The present study will measure the urea, H<sub>2</sub>O, and NH<sub>3</sub> permeabilities of mouse (m) UT-As expressed in *Lithobates catesbeianus* oocytes, a heterologous expression system developed and standardized by our laboratory. Briefly, urea uptake will be monitored using <sup>14</sup>C-urea, osmotic water permeability (Pf) will be computed using video microscopy after placing the oocytes in a hypotonic solution to monitor the rate of cell swelling, and NH<sub>3</sub> permeability will be measured using a microelectrode with a blunt tip to record the maximum transient change in pH at the surface of the oocyte (DpHS) caused by NH<sub>3</sub> influx. We believe that the results from this study will provide valuable insights into the role(s) of UTs-with their permeabilities to urea, water and NH<sub>3</sub> are an important nexus for integrating the excretion of nitrogenous wastes, water, and acid. Furthermore, deepening the knowledge of the mechanisms involved in the excretion of water and acid (ammonia) will improve our understanding of basic kidney function, and could potentially aid in the treatment of hydroelectrolytic and acid-base disorders.

**Start date:** 04-30-2019

**End date:** 10-30-2021

**Last modified:** 07-01-2024

**Copyright information:**

The above plan creator(s) have agreed that others may use as much of the text of this plan as they would like in their own plans, and customize it as necessary. You do not need to credit the creator(s) as the source of the language used, but using any of the plan's text does not imply that the creator(s) endorse, or have any relationship to, your project or proposal

---

# **Molecular and functional study of the movement of urea, water and ammonia (NH<sub>3</sub>/NH<sub>4</sub><sup>+</sup>) through the urea transporters expressed in *Lithobates catesbeianus* oocytes**

## **Data Collection**

---

### **What data will you collect or create?**

We have been doing surface protein expression and collecting measurements of urea, H<sub>2</sub>O, and NH<sub>3</sub> transport of mouse (m) UT-As expressed in *Lithobates catesbeianus* oocytes, a heterologous expression system developed and standardized by our laboratory.

### **How will the data be collected or created?**

The surface expression has been assessed by biotinylation and western blot. The Urea uptake was monitored using <sup>14</sup>C-urea. Osmotic water permeability (Pf) has been computed using video microscopy after placing the oocytes in a hypotonic solution to monitor the rate of cell swelling. The index of NH<sub>3</sub> permeability has been measured using a microelectrode with a blunt tip to record the maximum transient change in pH at the surface (s) of the oocyte (delta pH<sub>S</sub>) caused by NH<sub>3</sub> influx.

## **Documentation and Metadata**

---

### **What documentation and metadata will accompany the data?**

For surface expression we present western blots.

For the Urea uptake (C14 uptake), we create bar graphs demonstrating the urea uptake for each expressed UT.

For the Osmotic Water Permeability (Pf), we create bar graphs demonstrating the Pf for each expressed UT.

For the Index of NH<sub>3</sub> Permeability (Pf), we create bar graphs demonstrating the delta pH<sub>S</sub> for each expressed UT.

We perform statistical analysis to determine significant differences between groups.

## **Ethics and Legal Compliance**

---

### **How will you manage any ethical issues?**

All surgical and experimental procedures involving animals were previously approved by the Committee of Animal Care and Use at the Institute of Biomedical Sciences of the University of Sao Paulo (protocol # 7971160519).

### **How will you manage copyright and Intellectual Property Rights (IP/IPR) issues?**

All copyright materials will subject to negotiations with the publisher. Concerning intellectual property rights, these will shared among the Laboratory, University of Sao Paulo and Fapesp.

## **Storage and Backup**

---

### **How will the data be stored and backed up during the research?**

At the moment, we have sufficient storage for the data being collected. Also, we routinely save all our data in the computers in our laboratory and back them up on external hard drive (Samsung). The students of the lab - mentioned as data manager - have been responsible for backing up and recovering the data. In the event of an incident, the data will be recovered from the device that was not affected.

### **How will you manage access and security?**

I, the Principal Investigator of the laboratory, will manage access and security of the data. Due to the fact that our data are stored in two separate locations, not connected to each other, we feel that the risk of a security treat is minimal.

## **Selection and Preservation**

---

### **Which data are of long-term value and should be retained, shared, and/or preserved?**

Since each of our project is an extension of the previous project, all of our data is considered to have long term value.

### **What is the long-term preservation plan for the dataset?**

Our long term is to save all our data to a cloud server that we will purchase.

## **Data Sharing**

---

### **How will you share the data?**

Once all our data are in the cloud, anyone with permission will be able to access them.

### **Are any restrictions on data sharing required?**

All data will be shared with the people working in the laboratory and established collaborators.

## **Responsibilities and Resources**

---

### **Who will be responsible for data management?**

I, Raif Musa Aziz (the Principal Investigator), will be responsible.

**What resources will you require to deliver your plan?**

May be necessary to purchase some cloud storage spaces, but not at the moment.

---